

An Experimental Programme for the Collection and Use of Retouching Tools Made on Diaphyseal Bone Splinters by Millán Mozota Holgueras (ES)

Published by **EXARC** at <http://journal.exarc.net/issue-2013-2/ea/experimental-programme-collection-and-use-retouching-tools-made-diaphyseal-bone-splinters>

TABLE 3. Blank size and morphometry				
	Lenght (mm)	Width (mm)	Cortical width (mm)	Thickness (mm)
<i>Bos</i> fresh bone				
Mean	88,30	33,89	48,13	12
Max.	185	51,89	115	22,15
Min.	53,67	17,36	20	5,55
Median	79,735	32,57	47	11,72
Std. desv.	29,60	8,17	18,46	3,52
<i>Bos</i> dry bone				
Mean	107,52	39,54	52,2	12,64
Max.	161	61,29	87	27,74
Min.	56,41	22,68	23	5,43
Median	106,78	37,07	52	11,28
Std. desv.	29,151	10,63	17,87	5,61
<i>Cervus</i> Fresh bone				
Mean	89,23	18,70	32,01	5,60
Max.	130,98	29,31	86	9,1
Min.	53,9	11,07	17	3,26
Median	91,79	18,27	31	5,54
Std. desv.	18,79	3,23	10,74	1,02